

The Additive Completion of k th Powers

Wenguang Zhai

*Department of Mathematics, Shandong Normal University, Jinan 250014,
Shandong, People's Republic of China*

E-mail: wgzhai@jn-public.sd.cninfo.net

Communicated by A. Hildebrand

Received January 2, 1997

Let $k \geq 2$ be a fixed integer. For positive integers $M \leq N$, let $S_k(M, N)$ denote the set of all sets $A \subset [0, M]$ such that, for all positive integers $n \leq N$, n can be written as $n = a + b^k$ with $a \in A$ and b a positive integer. Define $f_k(M, N) = \min\{|A| : A \in S_k(M, N)\}$. Given $\varepsilon > 0$, we prove that there exists a $\delta > 0$ such that for all sufficiently large N

$$f_k(\delta N, N) \geq (k - \varepsilon) N^{1-1/k}.$$

© 1999 Academic Press

1. INTRODUCTION AND RESULTS

Suppose $k \geq 2$ is a fixed integer. For positive integers $M \leq N$, let $S_k(M, N)$ denote the set of all sets $A \subset [0, M]$ such that every positive integer $n \leq N$ can be written as $n = a + b^k$ with $a \in A$ and b a positive integer. Define $f_k(M, N) = \min\{|A| : A \in S_k(M, N)\}$. Many authors considered the problem of obtaining a lower bound for $f_k(M, N)$ in the case $M = N$ (see [1–6]. The latest result is due to J. Cilleruelo [4], who showed that

$$f_k(N, N) \geq N^{1-1/k} \left\{ \frac{1}{\Gamma(2-1/k) \Gamma(1+1/k)} + o(1) \right\}. \quad (1)$$

The aim of this paper is to estimate $f_k(M, N)$ when M is small compared to N . Let $M_k = M_k(N)$ denote the smallest integer M such that $S_k(M, N)$ is non-empty. For example, if S is an integer such that $S \geq 4k$, then $M_k(S^k - 1) = S^k - (S - 1)^k - 1$ and $M_k(S^k + 1) = (S + 1)^k - S^k - 1$. In general, we have the following proposition.

PROPOSITION. *Let $B = \lfloor N^{1/k} \rfloor$. Then*

$$B^k - (B - 1)^k - 1 \leq M_k \leq (B + 1)^k - B^k - 1. \quad (2)$$

Our main result is the following

THEOREM 1. *Let $k \geq 2$ be an integer. There exist constants $\varepsilon_0 = \varepsilon_0(k) > 0$ and $N_0 = N_0(k) > 1$ such that, if $N \geq N_0$ and $3k^2 N^{-1/2k} \leq \varepsilon \leq \varepsilon_0$, then we have*

$$f_k(\delta N, N) \geq (k - \varepsilon) N^{1-1/k},$$

where $\delta = \delta(\varepsilon) = \varepsilon^2/(9k^3)$.

The referee kindly suggested the following question: What can be said about the upper bound of $f_k(M, N)$? This question can be answered by

THEOREM 2. *For all integers $k \geq 2$ and $M_k(N) \leq M \leq N$ we have*

$$f_k(M, N) \leq (B + 1)^k - B^k, \quad (3)$$

where $B = [N^{1/k}]$.

From Theorem 1 and Theorem 2 we immediately get

COROLLARY. *Let $k \geq 2$ be an integer. There exist constants $0 < \delta_k < 1$ and $N_k > 1$ such that, if $N \geq N_k$ and $M_k/N < \delta < \delta_k$, then*

$$(k - 3k^{3/2}\delta^{1/2}) N^{1-1/k} \leq f_k(\delta N, N) \leq k(1 + kN^{-1/k}) N^{1-1/k}. \quad (4)$$

In particular, if $\delta = \delta(N)$ satisfies $\delta(N) \rightarrow 0$ as $N \rightarrow \infty$, then

$$f_k(\delta N, N) = (k + o(1)) N^{1-1/k} \quad (N \rightarrow \infty). \quad (5)$$

It may be conjectured that the asymptotic formula (5) holds uniformly for $M_k \leq M \leq N$. However, our method works only when $M = o(N)$.

For large k , we have the following result.

THEOREM 3. *Suppose $0 < \delta < 1$ is a fixed real number. There exist constants $k_0 = k_0(\delta) > 2$ and $N_k(\delta) > 2$ such that if $N \geq N_k(\delta)$ and $k \geq k_0$, then*

$$f_k(\delta N, N) \geq C(\delta) \frac{k^A}{\log k} \left(1 - \frac{\log(1 + 1/\delta)}{\log k} \right) N^{1-1/k},$$

where

$$C(\delta) = \frac{\delta \Delta^A \log(1 + 1/\delta)}{(1 + \Delta)^{1+\Delta}}, \quad \Delta = \frac{\log 1/\delta}{\log(1 + 1/\delta)}.$$

Remark. Our theorems also hold if b^k is replaced by $b^k + P_{k-1}(b)$ or $[b^c]$, where $P_{k-1}(b)$ is a polynomial of degree $k-1$ and $c \geq 2$ is any fixed real number.

2. PROOF OF THE PROPOSITION

The proof is short and simple. Let $A_1 = \{0, 1, 2, \dots, (B+1)^k - B^k - 1\}$, where $B = [N^{1/k}]$. For each $n \leq N$, there exists a positive integer b such that $b^k \leq n \leq (b+1)^k$. Put $a = n - b^k$. Then $n = a + b^k$ and $a \in A_1$. This implies that $S_k((B+1)^k - B^k - 1, N)$ is non-empty. Thus $M_k(N) \leq (B+1)^k - B^k - 1$.

Now let A_2 denote a set in $S_k(M_k, N)$. Consider the number $n = B^k - 1$. It is obvious that if we want to write n in the form $a + b^k$ with $a \in A_2$, we must have $b \leq B-1$ and $a \geq B^k - (B-1)^k - 1$. Thus $M_k(N) \geq B^k - (B-1)^k - 1$.

This completes the proof of the proposition.

3. PROOF OF THEOREM 1

Suppose that ε satisfies the condition of Theorem 1 and let $\delta = \varepsilon^2/(9k^3)$. Let $A \in S_k(\delta N, N)$. We consider the identity

$$\sum_{t \in A} x^t \sum_{b^k \leq N} x^{b^k} = \sum_{n \leq L} c(n) x^n, \quad (6)$$

where $L = \max\{n \mid n = a + b^k, a \in A, b^k \leq N\}$ and $c(n) = \#\{n = a + b^k \mid a \in A, b^k \leq N\}$. By the definition of $S_k(M, N)$ we have $c(n) \geq 1$ for all $n \leq N$ and $N \leq L \leq N + \delta N$.

Put $F(x) = \sum_{t \in A} x^t$, $G(x) = \sum_{b^k \leq N} x^{b^k}$ and $H(x) = \sum_{n \leq L} c(n) x^n$. Let $d = [2(1 - 1/k)/\varepsilon]$. By the Leibniz formula we get that

$$H^{(d)}(x) = \sum_{j=0}^d \binom{d}{j} F^{(j)}(x) G^{(d-j)}(x).$$

Setting $x = 1$, we obtain

$$H^{(d)}(1) = \sum_{j=0}^d \binom{d}{j} F^{(j)}(1) G^{(d-j)}(1). \quad (7)$$

We now evaluate $H^{(d)}(1)$ and $F^{(j)}(1)$, $G^{(j)}(1)$ for $0 \leq j \leq d$. Obviously we have

$$\begin{aligned} H^{(d)}(1) &= \sum_{d \leq n \leq L} c(n) n(n-1) \cdots (n-d+1) \\ &\geq \sum_{d \leq n \leq N} n(n-1) \cdots (n-d+1). \end{aligned} \quad (8)$$

where we used the fact $c(n) \geq 1$ for all $n \leq N$.

Let $y = y(x) = \sum_{n=0}^N x^n$. Then $(x-1)y = x^{N+1} - 1$. Using the Leibniz formula again we get

$$\begin{aligned} (x-1)y^{(d+1)}(x) + (d+1)y^{(d)}(x) \\ = (N+1)N(N-1) \cdots (N-d+1)x^{N-d}. \end{aligned}$$

Setting $x=1$ we obtain

$$y^{(d)}(1) = \frac{(N+1)N(N-1) \cdots (N-d+1)}{d+1}. \quad (9)$$

On the other hand,

$$y^{(d)}(1) = \sum_{d \leq n \leq N} n(n-1) \cdots (n-d+1). \quad (10)$$

Thus

$$\begin{aligned} H^{(d)}(1) &\geq \frac{(N+1)N(N-1) \cdots (N-d+1)}{d+1} \\ &> \frac{N^{d+1}}{d+1} e^{\sum_{j=1}^{d-1} \log(1-j/N)} \\ &= \frac{N^{d+1}}{d+1} e^{\sum_{j=1}^{d-1} -j/N + O(d^3/N^2)} \\ &= \frac{N^{d+1}}{d+1} e^{-(d(d-1)/2N) + O(d^3/N^2)} \\ &> \frac{N^{d+1}}{d+1} \left(1 - \frac{d(d-1)}{2N} + O(d^3/N^2) \right). \end{aligned} \quad (11)$$

To estimate $F^{(j)}(1)$ we observe that

$$F^{(0)}(1) = \sum_{t \in A} 1 = |A| \quad (12)$$

and

$$F^{(j)}(1) = \sum_{t \in A, t \geq j} t(t-1) \cdots (t-j+1) \leq |A| \delta^j N^j, \quad 1 \leq j \leq d. \quad (13)$$

Finally, we note that

$$G^{(0)}(1) = \sum_{b^k \leq N} 1 \leq N^{1/k} + 1 \quad (14)$$

and for $1 \leq j \leq d$

$$\begin{aligned} G^{(j)}(1) &= \sum_{j \leq b^k \leq N} b^k (b^k - 1) \cdots (b^k - j + 1) \\ &\leq \sum_{b^k \leq N} b^{jk} \leq \frac{N^{j+1/k}}{jk+1} + N^j. \end{aligned} \quad (15)$$

From (12)–(15) we obtain

$$\begin{aligned} H^{(d)}(1) &= \sum_{j=0}^d \binom{d}{j} F^{(j)}(1) G^{(d-j)}(1) \\ &\leq \sum_{j=0}^d \binom{d}{j} |A| \delta^j N^j \left(\frac{N^{d-j+1/k}}{(d-j)k+1} + N^{d-j} \right) \\ &\leq \frac{|A| N^{d+1/k}}{dk+1} \left(1 + (1+\delta)^d (dk+1) N^{-1/k} \right. \\ &\quad \left. + (dk+1) \sum_{j=1}^d \binom{d}{j} \delta^j \frac{1}{(d-j)k+1} \right) \\ &\leq \frac{|A| N^{d+1/k}}{dk+1} \left(1 + 2(dk+1) N^{-1/k} \right. \\ &\quad \left. + (dk+1) \sum_{j=1}^d \binom{d}{j} \delta^j \frac{1}{(d-j)k+1} \right), \end{aligned} \quad (16)$$

since

$$(1+\delta)^d < e^{d \log(1+\delta)} < e^{d\delta} < e^{1/2} < 2.$$

We now estimate the sum

$$\Sigma = \sum_{j=1}^d \binom{d}{j} \delta^j \frac{1}{(d-j)k+1}.$$

Write

$$\Sigma = \frac{d\delta}{(d-1)k+1} + \frac{d(d-1)\delta^2}{(d-2)k+1} + \sum_{3 \leq j \leq d/2} + \sum_{j > d/2},$$

say. We have

$$\sum_{j > d/2} < \delta^{d/2} \sum_{j > d/2} \binom{d}{j} < (2\delta^{1/2})^d < \frac{d\delta^2}{2k} \quad (17)$$

and

$$\begin{aligned} \sum_{3 \leq j \leq d/2} &< \frac{2}{dk} \sum_{3 \leq j \leq d} \binom{d}{j} \delta^j \\ &\leq \frac{2}{dk} ((1+\delta)^d - 1 - d\delta) = \frac{2}{dk} (e^{d \log(1+\delta)} - 1 - d\delta) \\ &< \frac{2}{dk} (e^{d\delta} - 1 - d\delta) < \frac{2d\delta^2}{k}, \end{aligned} \quad (18)$$

where we have used the inequalities $\log(1+x) < x$ ($x > 0$) and $e^x < 1+x+x^2$ ($0 < x < 1/2$).

Thus,

$$\Sigma < \frac{d\delta}{(d-1)k+1} + \frac{4d\delta^2}{k}, \quad (19)$$

if we notice

$$\frac{d(d-1)\delta^2}{(d-2)k+1} < \frac{3d\delta^2}{2k}.$$

Inserting (19) into (16) we get

$$H^{(d)}(1) < \frac{|A| N^{d+1/k}}{dk+1} \Delta_0 \quad (20)$$

with

$$\Delta_0 = 1 + 2(dk+1) N^{-1/k} + \frac{d(dk+1)\delta}{(d-1)k+1} + \frac{4d(dk+1)\delta^2}{k}.$$

Now from (11) and (20) we get

$$|A| > \Delta_1 N^{1-1/k}, \quad (21)$$

where

$$\begin{aligned}
 A_1 &= \frac{dk+1}{d+1} \left(1 - \frac{d(d-1)}{2N} + O\left(\frac{d^3}{N^2}\right) \right) A_0^{-1} \\
 &= k \left(1 - \frac{1-1/k}{d+1} \right) \left(1 - \frac{d(d-1)}{2N} + O\left(\frac{d^3}{N^2}\right) \right) A_0^{-1} \\
 &> k \left(1 - \frac{1-1/k}{d+1} \right) \left(1 - \frac{d(d-1)}{2N} + O\left(\frac{d^3}{N^2}\right) \right) \\
 &\quad \times \left(1 - 2(dk+1) N^{-1/k} - \frac{d(dk+1) \delta}{(d-1)k+1} - \frac{4d(dk+1) \delta^2}{k} \right) \\
 &\geq k - \varepsilon,
 \end{aligned}$$

if we recall the conditions on ε and the definitions of d and δ , and estimate A_0^{-1} using the inequality $1/(1+x) > 1-x$ ($|x| < 1$).

This completes the proof of Theorem 1.

4. PROOF OF THEOREM 2 AND COROLLARY

Without loss of generality, we may suppose $M \geq (B+1)^k - B^k - 1$; otherwise Theorem 2 and the corollary follow from the proposition.

Let $A_0 \in S_k(M, N)$ be such that $|A_0| = f_k(M, N)$. Since

$$\{0, 1, 2, \dots, (B+1)^k - B^k - 1\} \in S_k(M, N),$$

we have

$$|A_0| \leq (B+1)^k - B^k.$$

Thus (3) holds.

Now suppose $\delta_k > 0$ is a sufficiently small positive constant and suppose δ satisfies the conditions of Theorem 2. Set $\varepsilon = 3k^{3/2}\delta^{1/2}$. By Theorem 1 we have

$$|A_0| \geq (k - \varepsilon) N^{1-1/k} \tag{22}$$

if $M \leq \delta N$.

As for the upper bound we have

$$\begin{aligned}
 (B+1)^k - B^k &= k \int_B^{B+1} u^{k-1} du \\
 &= kB^{k-1} + k(k-1) \int_B^{B+1} du \int_B^u v^{k-2} dv \\
 &\leq kB^{k-1} + k(k-1)(B+1)^{k-2} \int_B^{B+1} du \int_B^u dv \\
 &\leq kB^{k-1} + k(k-1)(B+1)^{k-2} \leq kB^{k-1} + (kB)^2 \\
 &\leq kN^{1-1/k} + k^2 N^{1-2/k}.
 \end{aligned} \tag{23}$$

Now the corollary follows from (22) and (23).

5. PROOF OF THEOREM 3

Suppose $0 < \delta < 1$ is a fixed real number not depending on k and $A \in S_k(M, N)$. As in the proof of Theorem 1 we get

$$\begin{aligned}
 &\frac{(N+1)N(N-1)\cdots(N-d+1)}{d+1} \\
 &\leq |A| N^{d+1/k} \sum_{j=0}^d \binom{d}{j} \delta^j \frac{1}{(d-j)k+1}.
 \end{aligned} \tag{24}$$

We have

$$\begin{aligned}
 \sum_{j=0}^d \binom{d}{j} \delta^j \frac{1}{(d-j)k+1} &\leq \delta^d + k^{-1} \sum_{j=1}^d \binom{d}{j} \delta^j \\
 &\leq \delta^d + k^{-1}(1+\delta)^d \\
 &= \delta^d(1+k^{-1}(1+1/\delta)^d),
 \end{aligned} \tag{25}$$

provided $k > 2(1+1/\delta)^d$.

Thus we get

$$|A| N^{1/k-1} \geq \frac{\delta^{-d}}{d+1} (1-k^{-1}(1+1/\delta)^d) \left(1 - \frac{1}{N}\right) \cdots \left(1 - \frac{d-1}{N}\right). \tag{26}$$

Now take $d = \lceil (\log k/D)/\log(1+1/\delta) \rceil$, $D = 1 + A/A$, where A is defined in Section 1. Thus Theorem 3 follows from (26). It should be noted that the choice of D is best possible.

ACKNOWLEDGMENT

The author deeply thanks the referee for many helpful suggestions that improved the paper greatly.

REFERENCES

1. H. L. Abbott, On an additive completion of sets of integers, *J. Number Theory* **17** (1983), 135–143.
2. R. Balasubramanian, On the additive completion of squares, *J. Number Theory* **29** (1988), 10–12.
3. R. Balasubramanian and K. Soundararajan, On the additive completion of squares, II, *J. Number Theory* **40** (1992), 127–129.
4. J. Cilleruelo, The additive completion of k th-powers, *J. Number Theory* **44** (1993), 237–243.
5. R. Donagi and M. Herzog, On the additive completion of polynomial sets of integers, *J. Number Theory* **3** (1971), 150–154.
6. L. Moser, On an additive completion of sets of integers, in “Proc. Sympos. Pure Math.,” Vol. 8, Amer. Math. Soc., Providence, 1965.